





HORSE PROTOCOL SHEETS

SHOULDER PROTOCOL

For left shoulder. After doing steps on left shoulder, repeat first three steps on opposite side once. If both are painful, apply all steps to other side.




Large X: 20 seconds



Stars: 10 seconds each.


Do multiple stars all directions on various planes. Repeat 2-3 times.



Connect both sides: 10 seconds each. Finish with 20 minutes CES


KNEE/CARPUS PROTOCOL

For left knee. After doing steps on left knee, repeat first three steps on opposite side once. If both are painful, apply all steps to other side.




Large X: 20 seconds

Stars: 10 seconds each. Do multiple stars all directions on various planes. Repeat 2-3 times.



Probes are places directly across each other on various planes. 10 seconds each.




Connect both sides: 10 seconds each. Finish with 20 minutes CES


STIFLE PROTOCOL

For left stifle. Repeat first three steps on opposite leg once.


If both are painful, apply all steps to other side.




Large X: 20 seconds




Stars: 10 seconds each. Do multiple stars all directions. Repeat 2-3 times.



One probe inside the stifle and one probe on the outside.




Probes are places directly across each other on various planes. 10 seconds each.



Connect both sides: 10 seconds each. Finish with 20 minutes CES

HOCK PROTOCOL

For left hock. If both are painful, apply steps to other side.



Large X: 20 seconds

Stars: 10 seconds each.

Do multiple stars all directions.


Repeat 2-3 times.

Probes are places directly across each other on various planes. 10 seconds each.


Connect both sides: 10 seconds each. Finish with 20 minutes CES

Have a helper flex the joint. Probes opposite for 10 seconds.


SPINAL PAIN OR PARESIS PROTOCOL




Large X: 20 seconds



Multiple stars in various planes: 10 seconds.




Focus additional stars around area of greatest concern. Vary distance off center and the transmission from side to side.




Connect both sides: 10 seconds each. Finish with 20 minutes CES

CUSTOMIZED PROTOCOL

Draw lines on the figures for any specific zone needing customized microcurrent therapy. Use our 5-Step Protocol as a foundation for the treatment recommendations.





Finish with 20 minutes CES